AMOOR- I (2004 Course): SUMMER - 2016

Subject : Mathematical Foundations

Day : Monday
Date : 13/06/2016

S.D.E.

Time: 10.00 AM TO 1.00 PM Max Marks: 80 Total Pages: 2

N.B.:

- 1) Attempt **ANY THREE** questions from Section I and attempt **ANY TWO** questions from Section II.
- 2) Answers to both the sections should be written in the **SAME** answer book.
- 3) Figures to the right indicate **FULL** marks.

SECTION - I

Q.1 a) If U is universal set and

[08]

 $U = \{x \mid x \in \mathbb{N}, x \le 14\}, A = \{2, 4, 6, 8, 10, 12\},\$ $B = \{1, 3, 5, 7, 9, 10, 11, 12, 13\},$

find: i) n $(A \cup B)$

ii) $n(A \cup B)$ iii) $n(A' \cup B')$ iv) $n(A \cup A')$

b) If $f(x) = x^2 - 2x + 3$ and g(x) = 2x - 3, find: **i)** fog **ii)** fog⁻¹.

[08]

Q.2 a) Write truth tables for at least four logical connectives.

[08]

- **b)** Check whether the three points A(1, -2, 3), B(2, 3, -4) and C(0, -7, 10) are **[08]** collinear by using vectors.
- Q.3 a) Write a note on Eucledian algorithm. Explain it with one example.

[08]

- **b)** Find the inverse of the matrix $A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$. [08]
- Q.4 a) Write a short note on: Complexity of algorithm.

[08]

- **b)** If A = {2, 3, 4, 5, 6}, determine the truth value of each of the following [08] statements:
 - i) $\exists x \in A$, such that x + 3 = 10
 - ii) $\forall x \in A, x + 2 < 11$
 - iii) $\exists x \in A$, such that x + 2 < 5
 - iv) $\forall x \in A, x + 3 \ge 9$
- Q.5 a) By using mathematical induction, prove that $1+2+3+....+n=\frac{n(n+1)}{2}$. [08]
 - b) Explain ordered and unordered partitions.

[08]

P.T.O.

30132

SECTION - II

- Q.6 a) Find the number of permutations of the following words:

 i) MICROSOFT

 ii) CONSULTANCY.

 [08]
 - b) Find the value of x if: $4\log_2 16 + 3x\log_3 81 12\log_5 625 = 0$. [08]
- Q.7 a) If $A = \begin{bmatrix} 2 & 3 \\ 4 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & 6 \\ 2 & 3 \end{bmatrix}$ then find: i) AB ii) (AB)⁻¹
 - b) The probability that A can solve the problem is $\frac{3}{5}$ and that can be solved by B [08] is $\frac{2}{7}$. If they solve the problem independently, what is the probability that:
 - i) none solves the problem?
 - ii) only B solves the problem?
 - iii) the problem is solved?
 - iv) both of them solve the problem?
- Q.8 Write short notes on ANY FOUR of the following: [16]
 - a) Counting principles and its applications
 - b) Binomial distribution
 - c) Types of matrices
 - **d)** Recursively defined functions
 - e) Venn Diagram

* * * *